

Bayesian meta-analysis

What is a meta-analysis

“An analysis of analyses”

⇒ a single quantitative summary of studies answering the *same research question*

Ex: medical therapies effects are often evaluated in multiple different studies.

What is a meta-analysis

“An analysis of analyses”

⇒ a single quantitative summary of studies answering the *same research question*

Ex: medical therapies effects are often evaluated in multiple different studies.

⇒ pool individual observations from multiple studies ?

What is a meta-analysis

“An analysis of analyses”

⇒ a single quantitative summary of studies answering the *same research question*

Ex: medical therapies effects are often evaluated in multiple different studies.

⇒ pool individual observations from multiple studies ?

- ⚠ potential differences in the pooled experiments
- ⚠ only aggregated summary statistics estimates (“effect sizes”) available – alongside uncertainty (e.g. standard errors)

Study Heterogeneity

⚠ variations of the observed effects. . .

Study Heterogeneity

- ⚠ variations of the observed effects. . .
- within-study uncertainty, or
 - real heterogeneity in effect size between the different studies

Study Heterogeneity

- ⚠ variations of the observed effects. . .
- within-study uncertainty, or
 - real heterogeneity in effect size between the different studies

Often, different studies used different populations

⇒ potential extra-variability

+ different sample sizes ⇒ also impact the estimate and its variability

Meta-analysis random effects model

Common approach for meta-analysis:

$$y_i \sim \mathcal{N}(\theta_i, \sigma_i^2)$$

$$\theta_i \sim \mathcal{N}(\mu, \tau^2)$$

Meta-analysis random effects model

Common approach for meta-analysis:

$$y_i \sim \mathcal{N}(\theta_i, \sigma_i^2)$$

$$\theta_i \sim \mathcal{N}(\mu, \tau^2)$$

Hierarchical generalization of the fixed effect model:

$$y_i \sim \mathcal{N}(\mu, \sigma_i^2)$$

Meta-analysis random effects model

Common approach for meta-analysis:

$$y_i \sim \mathcal{N}(\theta_i, \sigma_i^2)$$

$$\theta_i \sim \mathcal{N}(\mu, \tau^2)$$

⇒ between study variability: $y_i \sim \mathcal{N}(\mu, \sigma_i^2 + \tau^2)$

Hierarchical generalization of the fixed effect model:

$$y_i \sim \mathcal{N}(\mu, \sigma_i^2)$$

⇒ assume same average effect for each study

Bayesian meta-analysis in practice

Meta-analysis: a perfect usecase for Bayesian analysis ?

Bayesian meta-analysis in practice

Meta-analysis: a perfect usecase for Bayesian analysis ?

- few observations
- informative *prior*
- sequential

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature

Going further

Scientific literature search

- ⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!
- ⚠ effect size estimate (along with their standard errors) must often be **transformed before** the meta-analysis

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!

⚠ effect size estimate (along with their standard errors) must often be **transformed before** the meta-analysis

Evidence synthesis

Meta-analysis \in evidence synthesis

e.g. meta-regression, mechanistic modeling, ...

Going further

Scientific literature search

- ⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!
- ⚠ effect size estimate (along with their standard errors) must often be **transformed before** the meta-analysis

Evidence synthesis

Meta-analysis \in evidence synthesis

e.g. meta-regression, mechanistic modeling, ...

Still active research domains:

- random effects model will down-weight studies with larger sample sizes
 - Serghiou & Goodman, *JAMA*, 2018

Going further

Scientific literature search

⚠ FIRST (!) exhaustive search of the scientific literature: hard !!!

⚠ effect size estimate (along with their standard errors) must often be **transformed before** the meta-analysis

Evidence synthesis

Meta-analysis \in evidence synthesis

e.g. meta-regression, mechanistic modeling, ...

Still active research domains:

- random effects model will down-weight studies with larger sample sizes
– Serghiou & Goodman, *JAMA*, 2018

a bug or a feature ?

Your turn !



Read ND Crins *et al.* Interleukin-2 Receptor Antagonists for Pediatric Liver Transplant Recipients: A Systematic Review and Meta-Analysis of Controlled Studies, *Pediatric Transplantation* 18(8):839, 2014. [DOI:10.1111/ptr.12362]

Practical: exercise 7